

Section[®] 2 EC

By WINFIELD

Grass weed control in soybeans, alfalfa and other crops

Postemergent Control for Actively Growing Grass Weeds

Section[®] 2 EC herbicide is applied postemergence for the control of a broad spectrum of actively growing annual and perennial grass weeds. It provides excellent control of volunteer cereal crops in broadleaf crops and of volunteer corn in Roundup Ready[®] soybeans when tank mixed with Cornerstone[®] or Roundup[®] brand herbicides. Section[®] 2 EC herbicide may be tank mixed with a variety of popular broadleaf herbicides and is rainfast within one hour.

Section[®] 2 EC herbicide may be used for weed control in alfalfa, beans, canola, cotton, flax, non-crop, ornamentals, peanuts, potatoes, soybeans, sugar beets and sunflower. It may also be used on a variety of fruits, vegetables and other broadleaf field crops.

For optimum weed control, always use an adjuvant such as Destiny[®] HC or Superb[®] HC with Section[®] 2 EC herbicide. Adjuvants help Section[®] 2 EC herbicide penetrate the leaf surface faster for more rapid weed control.

Ingredients

Clethodim	26.4%
Other ingredients	73.6%
TOTAL	100%

Section[®] 2 EC herbicide contains 2 pounds active ingredient per gallon.



Adjuvant Use

Section[®] 2 EC herbicide should be applied with crop oil concentrate, methylated seed oil, nonionic surfactant or UAN/AMS, depending on the specific crop.

Most field crops

1 pint per acre or 0.5% v/v of Superb[®] HC or Destiny[™] HC adjuvants. Do not use less than 1 pint of oil per acre by air. 1 to 2 quarts UAN or 2.5 to 4 pounds AMS may be added.

See label for vegetable crops, non-crop land and other uses.

To lock in spray coverage or deposition and lock out drift, include InterLock[®] adjuvant.

Packaging

| 4 x 1-gallon jugs

Section[®] 2 EC

By WINFIELD

Weeds Controlled

Section[®] 2 EC herbicide controls and/or suppresses a wide variety of annual and perennial grass weeds, including:

Annual Bromegrasses	Perennial bluegrass
Barnyardgrass	Quackgrass
Bermudagrass	Rabbitsfootgrass
Broadleaf signalgrass	Red rice
Canarygrass	Ryegrass (hardy, Italian)
Crabgrasses	Sandbur, field
Crowfootgrass	Shattercane
Fall panicum	Southwestern cupgrass
Fescue, tall	Sprangletop
Foxtail (giant, green, yellow)	Texas panicum
Foxtail barley	Volunteer cereals
Goosegrass	Volunteer corn
Itchgrass	Volunteer grain sorghum
Johnsongrass (rhizome, seedling)	Wild oats
Junglerice	Wild proso millet
Lovegrass	Wirestem muhly
Orchardgrass	Witchgrass
	Woolly cupgrass

Application Guidelines

Section[®] 2 EC herbicide offers a large window of application with optimal crop safety. Application should be made postemergence on actively growing grasses. Rates depend on weed species and size.

Section[®] 2 EC herbicide may be tank mixed with Cornerstone[®] or Roundup[®] brand herbicides and other glyphosate brands for control of volunteer corn.

Section[®] 2 EC herbicide may be applied by ground or air. It is compatible with MAX-IN[®] nutrients and many insecticides, such as Arctic[®], Grizzly[®] Z, Tundra[®], Tundra[®] Supreme and Yuma[®].

Weed Species	Weed Height (Inches)	Application Rate* (Ounces per Acre)
Annual Bromegrasses	2 to 6	6 to 8
Barnyardgrass	2 to 8	6 to 8
Crabgrasses	2 to 6**	6 to 8
Fall panicum	2 to 8	6 to 8
Fescue, tall	4 to 8*	8 to 16
Foxtail, giant	2 to 12	6 to 8
Foxtail, green	2 to 8	6 to 8
Foxtail, yellow	2 to 8	6 to 8
Johnsongrass, seedling	4 to 10	6 to 8
Quackgrass	4 to 12***	8 to 16
Ryegrass	2 to 6	6 to 8
Sandbur, field	2 to 6	6 to 8
Shattercane	6 to 18	6 to 8
Sprangletop	2 to 6	6 to 8
Volunteer cereals	2 to 6	6 to 8
Volunteer corn	4 to 12	4 to 6
Volunteer corn	12 to 24	6 to 8
Wild oats	2 to 6	6 to 8
Wild proso millet	2 to 10	6 to 8
Wirestem muhly	4 to 8***	8 to 16
Woolly cupgrass	2 to 8	6 to 8

*For many crops.

**Lateral growth.

***Perennials may require repeated applications.